

A remark of contraction semigroups on Banach spaces

Pei-Kee Lin

Department of Mathematics
Memphis State University
Memphis TN, 38152
USA

February 1, 2008

Abstract

Let X be a complex Banach space and let $J : X \rightarrow X^*$ be a duality section on X (i.e. $\langle x, J(x) \rangle = \|J(x)\| \|x\| = \|J(x)\|^2 = \|x\|^2$). For any unit vector x and any (C_0) contraction semigroup $T = \{e^{tA} : t \geq 0\}$, Goldstein proved that if X is a Hilbert space and if $|\langle T(t)x, J(x) \rangle| \rightarrow 1$ as $t \rightarrow \infty$, then x is an eigenvector of A corresponding to a purely imaginary eigenvalue. In this article, we prove the similar result holds if X is a strictly convex complex Banach space.

¹1991 Mathematics Subject Classification: 47B03, 47B15.

²Key words and phrases: eigenvalue, contraction, exposed point, strongly exposed point, semigroup.

Let X be a complex Banach space. Recall $x^* \in X^*$ is called *dual* of $x \in X$, if

$$\langle x, x^* \rangle = \|x^*\| \|x\| = \|x^*\|^2 = \|x\|^2.$$

a mapping $J : X \rightarrow X^*$ is said to be a *duality section* if $J(x)$ is dual to x for all $x \in X$. It is known that if X is a Hilbert space, then $J(x) = x$. In [3], Goldstein proved the following theorem.

Theorem 1 *Let A generate a (C_0) contraction semigroup $T = \{e^{tA} : t \geq 0\}$ on a complex Banach space. If X is a Hilbert space, then for any unit vector $x \in X$,*

$$\lim_{t \rightarrow \infty} |\langle T(t)x, J(x) \rangle| = 1$$

implies that $Ax = i\lambda x$ for some real number λ .

He also proved that Theorem 1 is not true if X replace by an L_∞ -space. In this article, we prove that Theorem 1 is true if and only if X is strictly convex.

Recall a Banach space is said to be *strictly convex* if for any two unit vectors $x, y \in X$, either $x = y$ or $\|x + y\| < 2$. By Hahn-Banach Theorem, X is strictly convex if and only if for any unit vectors x, y and any nonzero linear functional x^* , $\langle x, x^* \rangle = \|x^*\| = \langle y, x^* \rangle$ implies $x = y$.

Example 2 *Suppose x, y are two unit vectors on X and x^* is a linear functional on X such that*

$$\langle x, x^* \rangle = 1 = \|x^*\| = \langle y, x^* \rangle.$$

Let A be defined by

$$A(z) = i\langle z, x^* \rangle x + (i - \frac{1}{2})(z - \langle z, x^* \rangle x).$$

Then for any z ,

$$e^{tA}z = e^{it}\langle z, x^* \rangle x + e^{it}e^{-\frac{t}{2}}(z - \langle z, x^* \rangle x) = e^{it} \left(e^{-\frac{t}{2}}z + (1 - e^{-\frac{t}{2}})\langle z, x^* \rangle x \right).$$

Hence, A generates a (C_0) contraction semigroup $T = \{e^{tA} : t \geq 0\}$. On the other hand,

$$\langle T(t)y, x^* \rangle = e^{it} \langle \langle y, x^* \rangle x + e^{-\frac{t}{2}}(y - \langle y, x^* \rangle x), x^* \rangle = e^{it}$$

and

$$A(y) = ix + (i - \frac{1}{2})(y - x).$$

So Theorem 1 does not hold if X is not strictly convex.

For any complex number $z \neq 0$, let $\operatorname{sgn}(z) = \frac{\bar{z}}{|z|}$.

Theorem 3 *Let X be any strictly convex complex Banach space and let $J : X \rightarrow X^*$ be any duality section of X . If x is a unit vector and if $T = \{e^{tA} : t \geq 0\}$ is a (C_0) contraction semigroup generated by A such that*

$$\lim_{t \rightarrow \infty} |\langle T(t)x, J(x) \rangle| = 1,$$

then x is an eigenvector of A corresponding to a purely imaginary eigenvalue.

Proof. We claim that for any t , $T(t)x = c(t)x$ for some $|c(t)| = 1$.

Suppose it is not true. Then there is $t \geq 0$ such that $f(d) = \|x + dT(t)x\|$ is a continuous function from $\{d \in \mathbb{C} : |d| = 1\}$ into the interval $[0, 2)$. Hence, there is $\epsilon > 0$ such that

$$\|x + dT(t)x\| \leq 2 - \epsilon$$

for all $|d| = 1$. So if $s \geq 0$, then

$$\begin{aligned} 2 - \epsilon &\geq \left\| T(s) \left(x + \operatorname{sgn} \left(\frac{\langle T(s+t)x, J(x) \rangle}{\langle T(s)x, J(x) \rangle} \right) T(t)x \right) \right\| \\ &\geq \left| \left\langle T(s)x + \operatorname{sgn} \left(\frac{\langle T(s+t)x, J(x) \rangle}{\langle T(s)x, J(x) \rangle} \right) T(s+t)x, J(x) \right\rangle \right| \\ &= |\langle T(s)x, J(x) \rangle| + \left| \left\langle \operatorname{sgn} \left(\frac{\langle T(s+t)x, J(x) \rangle}{\langle T(s)x, J(x) \rangle} \right) T(s+t)x, J(x) \right\rangle \right|. \end{aligned}$$

This contradicts that

$$\lim_{s \rightarrow \infty} |\langle T(s)x, J(x) \rangle| = 1.$$

But

$$A(x) = \lim_{t \rightarrow 0^+} \frac{e^{tA}(x) - x}{t} = \lim_{t \rightarrow 0^+} \frac{(c(t) - c(0))x}{t} = c'(0)x.$$

This implies x is also an eigenvector of A corresponding to an eigenvalue $c'(0)$. Hence,

$$1 \geq \|e^A x\| = \|e^{c'(0)} x\| = |e^{c'(0)}|.$$

On the other hand,

$$\|e^A x\| \geq \lim_{t \rightarrow \infty} \|T(t)x\| \geq \lim_{t \rightarrow \infty} |\langle T(t)x, J(x) \rangle| = 1.$$

This implies $|e^{c'(0)}| = 1$, and so $c'(0)$ must be a purely imaginary number. $\square$

Let x be any unit vector in a complex Banach space such that if $y \neq x$ and $\|y\| = 1$, then $\|x+y\| < 2$. The proof above shows that if $\{T = e^{tA} : t \geq 0\}$ is a (C_0) contraction semigroup generated by A and if $\lim_{t \rightarrow \infty} |\langle T(t)x, J(x) \rangle| = 1$, then x is an eigenvector of A corresponding to a purely imaginary eigenvalue.

Let x^* be any unit vector in X^* , and let α be any real number between 0 and 1. The *slice* $S(B, x^*, \alpha)$ is the set

$$\{x : \|x\| \leq 1, \text{ and } \Re(\langle x, x^* \rangle) \geq 1 - \alpha\}.$$

Recall a unit vector x is called an *exposed point* (of the unit ball) if there is $x^* \in X^*$ such that for any $y \neq x$ with $\|y\| = 1$,

$$1 = \|x^*\| = \langle x, x^* \rangle > \Re\langle y, x^* \rangle.$$

The linear functional x^* is said to *expose* x . x is called a *strongly exposed point* if there is $x^* \in X^*$ such that $\langle x, x^* \rangle = 1$ and

$$\begin{aligned} 0 &= \lim_{\alpha \rightarrow 0^+} \text{diam } S(B, x^*, \alpha) \\ &= \lim_{\alpha \rightarrow 0^+} \sup\{\|y - z\| : y \in S(B, x^*, \alpha) \text{ and } z \in S(B, x^*, \alpha)\}. \end{aligned}$$

We say x^* *strongly exposes* x . One may ask the following question.

Question 1 *Let x be a unit vector. Suppose that x^* exposes x . Is x an eigenvector of A corresponding to a purely imaginary eigenvalue if $T = \{e^{tA} : t \geq 0\}$ is a (C_0) contraction semigroup generated by A such that*

$$\lim_{t \rightarrow \infty} |\langle T(t)x, x^* \rangle| = 1?$$

We do not know the general answer to this question. But the following theorems show that the answer is affirmative if x^* strongly exposes x or if X is a reflexive Banach space.

Theorem 4 *Let x be a unit vector of a complex Banach space X . Suppose that x^* strongly exposes x . If $T = \{e^{tA} : t \geq 0\}$ is a (C_0) contraction semigroup generated by A , and if*

$$\lim_{t \rightarrow \infty} |\langle T(t)x, x^* \rangle| = 1,$$

x is an eigenvector of A corresponding to a purely imaginary eigenvalue.

Proof. By the proof of Theorem 3, it enough to show that for each $s \geq 0$, x is an eigenvector of e^{sA} .

Since x^* strongly exposes x and

$$\lim_{t \rightarrow \infty} \operatorname{sgn}(\langle T(t)x, x^* \rangle) \langle T(t)x, x^* \rangle = 1,$$

$\operatorname{sgn}(\langle T(t)x, x^* \rangle)T(t)x$ converges to x . Hence, for any $s \geq 0$,

$$\begin{aligned} e^{sA}x &= e^{sA}(\lim_{t \rightarrow \infty} \operatorname{sgn}(\langle T(t)x, x^* \rangle)T(t)x) \\ &= \lim_{t \rightarrow \infty} \operatorname{sgn}(\langle T(t)x, x^* \rangle)T(s+t)x \\ &= \lim_{t \rightarrow \infty} \frac{\operatorname{sgn}(\langle T(t)x, x^* \rangle)}{\operatorname{sgn}(\langle T(s+t)x, x^* \rangle)}x. \end{aligned}$$

So x is an eigenvector of e^{sA} . We proved our theorem. $\square$

Theorem 5 *Let x be a unit vector of a complex reflexive Banach space X . Suppose that x^* exposes x . If $T = \{e^{tA} : t \geq 0\}$ is a (C_0) contraction semigroup generated by A , and if*

$$\lim_{t \rightarrow \infty} |\langle T(t)x, x^* \rangle| = 1,$$

x is an eigenvector of A corresponding to a purely imaginary eigenvalue.

Proof. As in the proof of Theorem 4, it is enough to show that x is an eigenvector of e^{sA} for all $s \geq 0$.

Fix $s \geq 0$ and a increasing positive sequence $\{t_n : n \in \mathbb{N}\}$ such that $\lim_{n \rightarrow \infty} t_n = \infty$. Since X is reflexive, by passing to subsequences of $\{e^{t_n A}x : n \in \mathbb{N}\}$ and $\{e^{(s+t_n)A}x : n \in \mathbb{N}\}$ we may assume that both of them converge weakly, say

$$\begin{aligned} y &= \text{w-} \lim_{n \rightarrow \infty} e^{t_n A}x \\ z &= \text{w-} \lim_{n \rightarrow \infty} e^{(s+t_n)A}x. \end{aligned}$$

But

$$\lim_{t \rightarrow \infty} |\langle e^{tA}x, x^* \rangle| = 1,$$

and x^* exposes x . So if $c = \langle y, x^* \rangle$, and $d = \langle z, x^* \rangle$, then $y = cx$, and $z = dx$. Hence

$$\begin{aligned} e^{sA}(cx) &= e^{sA}(\text{w-} \lim_{n \rightarrow \infty} e^{t_n A}x) \\ &= \text{w-} \lim_{n \rightarrow \infty} e^{sA}(e^{t_n A}x) \\ &= dx \end{aligned}$$

This implies x is an eigenvector of A corresponding to a purely imaginary eigenvalue. The proof is complete. $\square$

Remark 1. Let x be a nonzero vector, and let $T = \{T(t) = e^{tA} : t \geq 0\}$ be a (C_0) contraction semigroup generated by A in a complex Banach space X . Recently, Goldstein and Nagy [4] proved that if $|\langle T(t)x, x^* \rangle| \rightarrow |\langle x, x^* \rangle|$ as $t \rightarrow \infty$ for every $x^* \in X^*$, then x is an eigenvector of A corresponding to a purely imaginary eigenvalue.

References

- [1] J. Diestel, “*Geometry of Banach Space-selected topics*”, Lecture Notes in Math., Nr. 485, Springer-Verlag (1975).
- [2] J. A. Goldstein, *Semigroups of operators and applications*, Oxford University Press, New York, 1985.
- [3] J. A. Goldstein, *Extremal properties of contraction semigroups on Hilbert and Banach spaces* Bull. London Math. Soc. **25** (1993) 369-376.
- [4] J. A. Goldstein and B. Nagy, *An extremal property of contraction semigroups in Banach spaces*, preprint.